

Section 3.2 Properties of a Function's Graph

Objective 1: Determining the Intercepts of a Function

An **intercept** of a function is a point on the graph of a function where the graph either crosses or touches a coordinate axis. There are two types of intercepts:

- 1) The y -intercept, which is the y -coordinate of the point where the graph crosses or touches the y -axis.
- 2) The x -intercepts, which are the x -coordinates of the points where the graph crosses or touches the x -axis.

The y -intercept:


A function can have **at most** one y -intercept. The y -intercept exists if $x = 0$ is in the domain of the function. The y -intercept can be found by evaluating $f(0)$.

The x -intercept(s):

A function may have several (even infinitely many) x -intercepts. The x -intercepts, also called **real zeros**, can be found by finding all *real solutions* to the equation $f(x) = 0$. Although a function may have several zeros, only the real zeros are x -intercepts.

Objective 2: Determining the Domain and Range of a Function from its Graph

The domain of the graph below is the interval $[a, b)$ while the range is the interval $[c, d]$.




Objective 3: Determining Where a Function is Increasing, Decreasing or Constant

The graph of f rises from left to right on an open interval on which f is **increasing**. The values of $f(x)$ get larger as x gets larger on the interval.

The graph of f falls from left to right on an open interval in which f is **decreasing**. The values of $f(x)$ get smaller as x gets larger on the interval.

The graph of f is a horizontal line on an open interval in which f is **constant**. The values of $f(x)$ do not change as x gets larger on the interval.



The function shown above is increasing on the interval (c, d) .


The function shown above is decreasing on the interval (a, b) .

The function shown above is constant on the interval (b, c) .


Objective 4: Determining Relative Maximum and Relative Minimum Values of a Function

When a function changes from increasing to decreasing at a point $(c, f(c))$, then f is said to have a relative maximum at $x = c$. The relative maximum value is $f(c)$.

Similarly, when a function changes from decreasing to increasing at a point $(c, f(c))$, then f is said to have a relative minimum at $x = c$. The relative minimum value is $f(c)$.



The relative minimum occurs at $x = c$, the relative minimum value is $f(c)$.






The relative maximum occurs at $x = c$, the relative maximum value is $f(c)$.

The word “relative” indicates that the function obtains a maximum or minimum value relative to some open interval. It is not necessarily the maximum (or minimum) value of the function on the entire domain.






A relative maximum cannot occur at an endpoint and must occur in an open interval. This applies to a relative minimum as well.

Objective 5: Determining if a Function is Even, Odd or Neither



Definition: A function f is **even** if for every x in the domain, $f(-x) = f(x)$. Even functions are symmetric about the y -axis. For each point (x, y) on the graph, the point $(-x, y)$ is also on the graph.



Definition: A function f is **odd** if for every x in the domain, $f(-x) = -f(x)$. Odd functions are symmetric about the origin. For each point (x, y) on the graph, the point $(-x, -y)$ is also on the graph.

Objective 6: Determining Information about a Function from a Graph