

# WEAK MAPS AND THE TUTTE POLYNOMIAL

CHRISTINE CHO AND JAMES OXLEY

ABSTRACT. Let  $M$  and  $N$  be matroids such that  $N$  is the image of  $M$  under a rank-preserving weak map. Generalizing results of Lucas, we prove that, for  $x$  and  $y$  positive,  $T(M; x, y) \geq T(N; x, y)$  if and only if  $x + y \geq xy$  or  $M \cong N$ . We give a number of consequences of this result.

## 1. INTRODUCTION

Terminology and notation used here will follow [6] unless otherwise stated. Given two rank- $r$  matroids  $M$  and  $N$ , a bijective map from  $E(M)$  to  $E(N)$  is a *rank-preserving weak map* if every basis of  $N$  is the image of a basis of  $M$ . We will write  $M \xrightarrow{rp} N$  if  $N$  is a rank-preserving weak-map image of  $M$ .

The following theorem of Lucas [5] shows that the numbers of bases, independent sets, and spanning sets of  $M$  are greater than the corresponding numbers for  $N$  if there is a rank-preserving weak map from  $M$  to  $N$ . Note that  $T(M; 1, 1)$ ,  $T(M; 2, 1)$ , and  $T(M; 1, 2)$  count the numbers of bases, independent sets, and spanning sets of  $M$ , respectively, where  $T(M; x, y)$  is the Tutte polynomial of  $M$ .

**Theorem 1.** *For non-isomorphic matroids  $M$  and  $N$ , if there is a rank-preserving weak map from  $M$  to  $N$ , then*

- (i)  $T(M; 1, 1) > T(N; 1, 1)$ ;
- (ii)  $T(M; 2, 1) > T(N; 2, 1)$ ;
- (iii)  $T(M; 1, 2) > T(N; 1, 2)$ ;
- (iv)  $T(M; x, 0) \geq T(N; x, 0)$  for all  $x > 0$  unless  $M$  has a loop;
- (v)  $T(M; 0, y) \geq T(N; 0, y)$  for all  $y > 0$  unless  $M$  has a coloop.

The main result of the paper is the following generalization of the last theorem.

---

*Date:* January 21, 2025.

*2020 Mathematics Subject Classification.* 05B35.

*Key words and phrases.* matroid, weak map, Tutte polynomial.

**Theorem 2.** *Let  $x$  and  $y$  be positive real numbers. Let  $M$  and  $N$  be matroids such that there is a rank-preserving weak map from  $M$  to  $N$ . Then  $T(M; x, y) \geq T(N; x, y)$  if and only if  $x + y \geq xy$  or  $M \cong N$ .*

In a matroid  $M$ , an element  $f$  is *freer* than an element  $g$  if  $g$  is contained in the closure of every circuit containing  $f$ . As a consequence of Theorem 2, we deduce that if  $f$  is freer than  $g$  in a matroid  $M$ , then the numbers of bases, circuits, and hyperplanes of  $M$  containing  $f$  are at least as large as the corresponding numbers of sets containing  $g$ .

The next section presents some preliminaries. The main result is proved in Section 3. The last section contains consequences of the main theorem.

## 2. PRELIMINARIES

In a matroid  $M$ , the *nullity* of a subset  $X$  of  $E(M)$  is  $|X| - r_M(X)$ . For matroids  $M$  and  $N$ , a bijection  $\varphi : E(M) \rightarrow E(N)$  is a *weak map* if  $\varphi^{-1}(I) \in \mathcal{I}(M)$  whenever  $I \in \mathcal{I}(N)$ . If  $r(M) = r(N)$ , then  $\varphi$  is a rank-preserving weak map from  $M$  to  $N$ . Although it is not required that a weak map be bijective, we will only consider bijective weak maps. Such maps have the following attractive property (see, for example, [6, Corollary 7.3.13]).

**Lemma 3.** *If  $\varphi : M \rightarrow N$  is a rank-preserving weak map from  $M$  to  $N$ , then  $\varphi$  is a rank-preserving weak map from  $M^*$  to  $N^*$ .*

For a matroid  $M$  with ground set  $E$ , the *Tutte polynomial*  $T(M; x, y)$  of  $M$  is defined by

$$T(M; x, y) = \sum_{A \subseteq E} (x-1)^{r(M)-r(A)} (y-1)^{|A|-r(A)}.$$

It is well known that  $T(M^*; x, y) = T(M; y, x)$  for a matroid  $M$  and its dual  $M^*$ . An in-depth account of the Tutte polynomial and its applications can be found in [1].

Before proving the main result in general, we prove it in the specific case when  $N$  is comprised solely of loops and coloops.

**Lemma 4.** *Let  $x > 0$  and  $y > 0$ . Let  $M$  be a matroid with rank  $k$ , nullity  $m$ , and  $|E(M)| \geq 2$ . Then  $T(M; x, y) \geq x^k y^m$  if and only if  $x + y \geq xy$  or  $M \cong U_{k,k} \oplus U_{0,m}$ .*

*Proof.* Suppose  $x + y < xy$  and  $M \not\cong U_{k,k} \oplus U_{0,m}$ . To show that  $T(M; x, y) < x^k y^m$ , we argue by induction on  $|E(M)|$ . If  $|E(M)| = 2$ , then  $M \cong U_{1,2}$  and  $T(M; x, y) = x + y < xy$ , and the result holds. Now assume that the result holds for  $|E(M)| < n$  and let  $|E(M)| = n \geq 3$ .

Since  $M \not\cong U_{k,k} \oplus U_{0,m}$ , there is an element  $e$  of  $M$  that is neither a loop nor a coloop. Then  $T(M; x, y) = T(M \setminus e; x, y) + T(M/e; x, y)$ .

Suppose that  $M \setminus e \cong U_{k,k} \oplus U_{0,m-1}$ . Then  $M \cong U_{s,s+1} \oplus U_{k-s,k-s} \oplus U_{0,m-1}$  for some  $s \leq k$ . Thus

$$\begin{aligned} T(M; x, y) &= T(U_{s,s+1}; x, y)x^{k-s}y^{m-1} \\ &= (x^s + x^{s-1} + \cdots + x + y)x^{k-s}y^{m-1}. \end{aligned}$$

To show that  $T(M; x, y) < x^k y^m$ , it suffices to show that  $x^s + x^{s-1} + \cdots + x + y < x^s y$ . This is certainly true if  $s = 1$  as  $x + y < xy$ . Now suppose  $s \geq 2$ . Then, arguing by induction on  $s$ , we have

$$\begin{aligned} x^s + x^{s-1} + \cdots + x^2 + x + y &< x^s + x^{s-1} + \cdots + x^2 + xy \\ &= x(x^{s-1} + \cdots + x^2 + x + y) \\ &< x(x^{s-1}y) \\ &= x^s y. \end{aligned}$$

We conclude that  $T(M; x, y) < x^k y^m$  when  $M \setminus e \cong U_{k,k} \oplus U_{0,m-1}$ .

We may now assume that  $M \setminus e \not\cong U_{k,k} \oplus U_{0,m-1}$ . By duality, we may also assume that  $M/e \not\cong U_{k-1,k-1} \oplus U_{0,m}$ . Therefore, by induction on  $|E(M)|$ ,

$$\begin{aligned} T(M; x, y) &= T(M \setminus e; x, y) + T(M/e; x, y) \\ &< x^k y^{m-1} + x^{k-1} y^m \\ &= x^{k-1} y^{m-1} (x + y) \\ &< x^{k-1} y^{m-1} (xy) \\ &= x^k y^m, \end{aligned}$$

as desired.

Conversely, since  $T(M; x, y) = x^k y^m$  when  $M \cong U_{k,k} \oplus U_{0,m}$ , we may assume that  $M \not\cong U_{k,k} \oplus U_{0,m}$ . Now suppose  $x + y \geq xy$ . If  $|E(M)| = 2$ , then  $M \cong U_{1,2}$  and  $T(M; x, y) = x + y \geq xy$ . Thus the result holds if  $|E(M)| = 2$ .

Assume the result holds for  $|E(M)| < n$  and let  $|E(M)| = n \geq 3$ . As  $M \not\cong U_{k,k} \oplus U_{0,m}$ , it follows that  $M$  has an element  $e$  that is neither

a coloop nor a loop. By the induction assumption,

$$\begin{aligned}
T(M; x, y) &= T(M \setminus e; x, y) + T(M/e; x, y) \\
&\geq x^k y^{m-1} + x^{k-1} y^m \\
&= x^{k-1} y^{m-1} (x + y) \\
&\geq x^{k-1} y^{m-1} (xy) \\
&= x^k y^m.
\end{aligned}$$

We conclude that the lemma holds.  $\square$

### 3. PROOF OF THE MAIN THEOREM

The argument uses the sign function  $\text{sgn} : \mathbb{R} \rightarrow \{-1, 0, 1\}$  where

$$\text{sgn}(x) = \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ 1 & \text{if } x > 0. \end{cases}$$

*Proof of Theorem 2.* It suffices to prove the result when  $M$  and  $N$  have a common ground set  $E$  and the rank-preserving weak map from  $M$  to  $N$  is the identity map. We will argue by induction on  $|E|$  that  $\text{sgn}(T(M) - T(N)) = \text{sgn}(x + y - xy)$  whenever  $M \neq N$  where we have abbreviated  $T(M; x, y)$  as  $T(M)$ .

Let  $|E| = 2$ . Since  $U_{1,2}$  and  $U_{1,1} \oplus U_{0,1}$  are the only 2-element matroids of equal rank, we must have that  $M \cong U_{1,2}$  and  $N \cong U_{1,1} \oplus U_{0,1}$ . As  $T(U_{1,2}; x, y) = x + y$  and  $T(U_{1,1} \oplus U_{0,1}; x, y) = xy$ , we see that the result holds for  $|E| = 2$ .

Assume the result holds for  $|E| < n$  and let  $|E| = n \geq 3$ . Take  $e \in E$ . If  $e$  is a coloop of  $M$ , then  $e$  is a coloop of  $N$ , so  $T(M) = xT(M \setminus e)$  and  $T(N) = xT(N \setminus e)$ . Therefore, as  $x > 0$ ,

$$\text{sgn}(T(M) - T(N)) = \text{sgn}(T(M \setminus e) - T(N \setminus e)) = \text{sgn}(x + y - xy).$$

Applying a similar argument to the dual, we see that the assertion holds if  $M$  has a loop.

Suppose  $e$  is not a loop or a coloop of  $N$ . Then

$$\begin{aligned}
\text{sgn}(T(M) - T(N)) &= \text{sgn}(T(M \setminus e) + T(M/e) - T(N \setminus e) - T(N/e)) \\
&= \text{sgn}(T(M \setminus e) - T(N \setminus e) \\
&\quad + T(M/e) - T(N/e)). \tag{1}
\end{aligned}$$

Since  $M \neq N$ , we have that  $M \setminus e \neq N \setminus e$  or  $M/e \neq N/e$ .

Suppose that  $M \setminus e \neq N \setminus e$  and  $M/e \neq N/e$ . Then, by the induction assumption,

$$\operatorname{sgn}(T(M \setminus e) - T(N \setminus e)) = \operatorname{sgn}(x + y - xy) = \operatorname{sgn}(T(M/e) - T(N/e)).$$

Thus  $\operatorname{sgn}(T(M \setminus e) - T(N \setminus e) + T(M/e) - T(N/e)) = \operatorname{sgn}(x + y - xy)$ , that is,  $\operatorname{sgn}(T(M) - T(N)) = \operatorname{sgn}(x + y - xy)$ .

Now suppose that  $M \setminus e = N \setminus e$  or  $M/e = N/e$ . Then, by (1),

$$\begin{aligned} \operatorname{sgn}(T(M) - T(N)) &= \begin{cases} \operatorname{sgn}(T(M \setminus e) - T(N \setminus e)) & \text{if } M/e = N/e, \\ \operatorname{sgn}(T(M/e) - T(N/e)) & \text{if } M \setminus e = N \setminus e, \end{cases} \\ &= \operatorname{sgn}(x + y - xy) \end{aligned}$$

where the last step follows by the induction assumption.

Finally, if every element of  $N$  is a loop or a coloop, then  $N \cong U_{k,k} \oplus U_{0,m}$ , so  $T(N; x, y) = x^k y^m$ . Since  $M \xrightarrow{r_p} N$ , the matroid  $M$  has rank  $k$  and nullity  $m$ . The result follows immediately from Lemma 4.  $\square$

The significance of the equation  $x + y = xy$  was noted by Jaeger, Vertigan, and Welsh [4] in their analysis of the computational complexity of the Tutte polynomial. Evidently  $x + y = xy$  if and only if  $(x, y)$  is a point on the hyperbola  $H_1$  defined by the equation  $(x - 1)(y - 1) = 1$ . It is straightforward to prove that  $T(M; x, y) = (x - 1)^{r(M)} y^{|E|}$  for all  $(x, y) \in H_1$ . Therefore, for any two matroids  $M$  and  $N$  that have the same rank and the same ground set,  $T(M; x, y) = T(N; x, y)$  for all  $(x, y) \in H_1$ , that is, for all  $(x, y)$  for which  $x + y = xy$ .

#### 4. CONSEQUENCES

A flat  $F$  of a matroid  $M$  is *cyclic* if  $F$  is a union of circuits. Given distinct elements  $f$  and  $g$  of a matroid  $M$ , Duke [2] defined  $f$  to be freer than  $g$  in  $M$  if  $g$  is contained in every cyclic flat of  $M$  containing  $f$ . It is worth noting that, if  $f$  is a coloop of  $M$ , then  $f$  is vacuously freer than  $g$  for all  $g \in E(M) - \{f\}$ . Likewise, if  $g$  is a loop of  $M$ , then  $f$  is freer than  $g$  for all  $f \in E(M) - \{g\}$ . Consequently, our discussion of relative freedom is primarily concerned with elements of  $M$  that are neither loops nor coloops.

Duke showed that relative freedom extends nicely to both duals and minors. If  $f$  is freer than  $g$  in  $M$ , then  $g$  is freer than  $f$  in  $M^*$ . Moreover,  $f$  is freer than  $g$  in  $M \setminus X / Y$  for all disjoint subsets  $X$  and  $Y$  of  $E(M)$ .

This section explores the notion of relative freedom of elements of a matroid and its connection to weak maps and the Tutte polynomial.

The following result provides our first direct link between relative freedom and rank-preserving weak maps.

Define the map  $\varphi_{gf} : E(M/f) \rightarrow E(M/g)$  by taking  $\varphi_{gf}(g) = f$  and  $\varphi_{gf}(e) = e$  for all  $e \neq g$ .

**Lemma 5.** *If  $f$  is freer than  $g$  in a matroid  $M$  and  $g$  is not a loop of  $M$ , then  $\varphi_{gf}$  is a rank-preserving weak map from  $M/f$  to  $M/g$ .*

*Proof.* Let  $I$  be independent in  $M/g$ . Then  $I \cup g$  is independent in  $M$ . Suppose  $f \notin I$ . Then  $\varphi_{gf}^{-1}(I) = I$ . If  $I$  is dependent in  $M/f$ , then  $M$  has a circuit  $C$  such that  $C \subseteq I \cup f$ . Moreover,  $f \in C$  since  $I$  is independent in  $M$ . As  $f$  is freer than  $g$  in  $M$ , we see that  $g \in \text{cl}_M(C)$ . Then  $I \cup g$  contains a circuit of  $M$ , a contradiction. Therefore  $I$  is independent in  $M/f$ .

Suppose  $f \in I$ . Then  $f \in I \cup g$  and  $I \cup g$  is independent in  $M$ . Therefore  $\varphi_{gf}^{-1}(I)$ , which equals  $(I \cup g) - f$ , is independent in  $M/f$ .  $\square$

The next result follows immediately from Theorem 2 and Lemma 5. The straightforward proof is omitted.

**Corollary 6.** *Let  $x > 0$  and  $y > 0$ . If  $f$  is freer than  $g$  in  $M$  and  $g$  is not a loop of  $M$ , then  $T(M/f; x, y) \geq T(M/g; x, y)$  if and only if  $x + y \geq xy$  or  $M/f \cong M/g$ .*

**Corollary 7.** *Let  $x > 0$  and  $y > 0$ . If  $f$  is freer than  $g$  in  $M$  and  $g$  is not a coloop of  $M$ , then  $T(M \setminus f; x, y) \leq T(M \setminus g; x, y)$  if and only if  $x + y \geq xy$  or  $M \setminus f \cong M \setminus g$ .*

*Proof.* Since  $g$  is freer than  $f$  in  $M^*$ , it follows by Corollary 6 that  $T(M^*/f; y, x) \leq T(M^*/g; y, x)$  if and only if  $x + y \geq xy$  or  $M^*/f \cong M^*/g$ . Thus, by duality, we have  $T(M \setminus f; x, y) \leq T(M \setminus g; x, y)$  if and only if  $x + y \geq xy$  or  $M \setminus f \cong M \setminus g$ .  $\square$

The following result lists several consequences of Corollary 6. We use  $b(M)$ ,  $W_k(M)$ ,  $h(M)$ , and  $\gamma(M)$  to represent the numbers of bases, rank- $k$  flats, hyperplanes, and circuits of  $M$ , respectively. To specify the numbers of such sets containing some element  $e$  of  $M$ , we write, for example,  $b(e; M)$  and  $W_k(e; M)$ .

**Corollary 8.** *If  $f$  is freer than  $g$  in  $M$ , then*

- (i)  $b(f; M) \geq b(g; M)$ ;
- (ii)  $W_k(f; M) \geq W_k(g; M)$  for all  $k \geq 0$ , provided  $g$  is not a loop of  $M$ ;
- (iii)  $h(f; M) \geq h(g; M)$ , provided  $g$  is not a loop of  $M$ ;
- (iv)  $\gamma(f; M) \geq \gamma(g; M)$ , provided  $f$  is not a coloop of  $M$ .

*Proof.* For (i), note that  $b(e; M) = b(M/e)$  for  $e \in E(M)$  as long as  $e$  is not a loop. If  $g$  is a loop of  $M$ , then  $b(g; M) = 0$ , so (i) holds. Assume  $g$  is not a loop of  $M$ . As  $f$  is freer than  $g$  and  $g$  is not a loop,  $f$  is not a loop. Thus the map  $\varphi_{gf}$  is a rank-preserving weak map from  $M/g$  to  $M/f$ . By Theorem 2, we have  $b(M/f) = T(M/f; 1, 1) \geq T(M/g; 1, 1) = b(M/g)$ . Thus (i) holds.

To prove (ii), observe that, when  $e$  is not a loop of  $M$ , a set  $X$  is a rank- $k$  flat of  $M$  if and only if  $X - e$  is a rank- $(k - 1)$  flat of  $M/e$ . Suppose  $g$  is not a loop of  $M$ . Then, since  $\varphi_{gf}$  is a rank-preserving weak map from  $M/f$  to  $M/g$ , it follows by [7, Proposition 9.3.3], that  $W_{k-1}(M/f) \geq W_{k-1}(M/g)$  for all  $k \geq 1$ . Thus  $W_k(f; M) \geq W_k(g; M)$  for all  $k \geq 1$ . Also  $W_0(f; M) = 0 = W_0(g; M)$  since neither  $f$  nor  $g$  is a loop of  $M$ . Thus (ii) holds. Hence so does (iii).

For (iv), observe that  $\gamma(e; M) = h(M^*) - h(e; M^*)$ . Assume  $f$  is not a coloop of  $M$ . Since  $g$  is freer than  $f$  in  $M^*$ , we have, by (iii), that  $h(f; M^*) \leq h(g; M^*)$ . Therefore

$$h(M^*) - h(f; M^*) \geq h(M^*) - h(g; M^*)$$

and (iv) holds.  $\square$

Let  $\gamma'(e; M)$  be the number of spanning circuits of  $M$  containing an element  $e$  of  $M$ .

**Corollary 9.** *If  $f$  is freer than  $g$  in  $M$  and  $f$  is not a coloop of  $M$ , then  $\gamma'(f; M) \geq \gamma'(g; M)$ .*

*Proof.* Take a spanning circuit  $D$  of  $M$  containing  $g$  but not  $f$ . Then  $D = B \cup g$  for some basis  $B$  of  $M$ . Suppose  $B \cup f$  is not a circuit of  $M$ . Then  $B \cup f$  properly contains a circuit  $C$  of  $M$  and  $f \in C$ . Hence  $\text{cl}(C)$  contains  $g$ . The set  $C - f$  spans  $C$ , so  $g \in \text{cl}(C - f)$ . Thus  $(C - f) \cup g$  is a dependent set that is a proper subset of the circuit  $D$ , a contradiction.  $\square$

**Lemma 10.** *The following are equivalent for elements  $f$  and  $g$  in a matroid  $M$ .*

- (i)  $f$  is freer than  $g$  in  $M$ ;
- (ii)  $b(f; N) \geq b(g; N)$  for all restrictions  $N$  of  $M$  containing  $\{f, g\}$ .

*Proof.* Suppose (i) holds. Then  $f$  is freer than  $g$  in all restrictions  $N$  of  $M$  containing  $\{f, g\}$ , so (ii) holds by Corollary 8(i).

Suppose (ii) holds and suppose  $f$  is not freer than  $g$ . Then  $M$  has a cyclic flat  $F$  containing  $f$  and avoiding  $g$ . Let  $N' = M|(F \cup g)$ . Note that  $g$  is a coloop of  $N'$ . Then  $b(g; N') = b(N')$ . As  $b(f; N') \geq$

$b(g; N')$ , it follows that  $b(f; N') = b(N')$ . Thus  $f$  is a coloop of  $N'$ , a contradiction.  $\square$

In the remaining results of this paper, we investigate several instances of equality holding between the number of distinguished sets of  $M$  containing  $f$  and the number of such sets containing  $g$ . Let  $x$  and  $y$  be elements of  $M$ . Then  $x$  and  $y$  are *clones* in  $M$  if and only if the bijection from  $E(M)$  to  $E(M)$  that interchanges  $x$  and  $y$  and fixes every other element is an isomorphism. It was shown in [3, Proposition 4.9] that  $x$  and  $y$  are clones if and only if the set of cyclic flats containing  $x$  is equal to the set of cyclic flats containing  $y$ . Thus  $x$  and  $y$  are clones if and only if  $x$  is freer than  $y$ , and  $y$  is freer than  $x$  in  $M$ .

**Theorem 11.** *Let  $f$  be freer than  $g$  in  $M$ . Then  $b(f; M) = b(g; M)$  if and only if  $f$  and  $g$  are clones in  $M$ .*

*Proof.* If  $f$  and  $g$  are clones in  $M$ , then clearly  $b(f; M) = b(g; M)$ . To prove the converse, suppose  $b(f; M) = b(g; M)$ . First assume that  $g$  is a loop of  $M$ . Then  $b(g; M) = 0 = b(f; M)$ . Thus  $f$  is a loop of  $M$ . Therefore  $f$  and  $g$  are clones in  $M$ . Similarly, if  $g$  is a coloop of  $M$ , then  $f$  and  $g$  are clones in  $M$ . We may assume that  $f$  and  $g$  are neither loops nor coloops. Thus  $b(f; M) = b(M/f)$  and  $b(g; M) = b(M/g)$ .

Let  $|E(M)| \in \{2, 3\}$ . Since  $f$  and  $g$  are not loops or coloops in  $M$ , we have that  $M \in \{U_{1,2}, U_{2,3}, U_{1,3}, U_{1,2} \oplus U_{0,1}, U_{1,2} \oplus U_{1,1}\}$ . It is straightforward to check that, in these cases,  $f$  and  $g$  are clones.

Assume the result holds for  $|E(M)| < n$  and let  $|E(M)| = n \geq 4$ . Suppose  $f$  and  $g$  are not clones in  $M$ . Take an element  $e \in E(M) - \{f, g\}$ . If  $e$  is a loop or a coloop in  $M$ , then  $b(f; M \setminus e) = b(f; M)$  and  $b(g; M \setminus e) = b(g; M)$ . Thus  $b(f; M \setminus e) = b(g; M \setminus e)$ . By the induction assumption,  $f$  and  $g$  are clones in  $M \setminus e$ . Hence  $f$  and  $g$  are clones in  $M$ , a contradiction. Thus  $e$  is neither a loop nor a coloop of  $M$ .

Suppose  $\{e, f\}$  is a circuit of  $M$ . Then  $\{e, f, g\}$  is contained in a parallel class since  $f$  is freer than  $g$  and  $g$  is not a loop in  $M$ , so  $f$  and  $g$  are clones of  $M$ . Thus we may assume that  $e$  is not a loop of  $M/f$ . If  $\{e, g\}$  is a circuit of  $M$ , then  $e$  is a loop in  $M/g$ , so  $b(M/g) = b(M/g \setminus e)$ . Therefore  $b(M/f \setminus e) + b(M/f/e) = b(M/g \setminus e)$ . By Lemma 10,  $b(M/f \setminus e) \geq b(M/g \setminus e)$ , so  $b(M/f) > b(M/g)$ , a contradiction.

Now  $e$  is not a loop or a coloop of  $M$ ,  $M/f$ , or  $M/g$  and it follows that  $b(M/f) = b(M/f \setminus e) + b(M/f/e)$  and  $b(M/g) = b(M/g \setminus e) + b(M/g/e)$ . By assumption,

$$b(M/f \setminus e) + b(M/f/e) = b(M/g \setminus e) + b(M/g/e).$$

By Lemma 10,  $b(M/f \setminus e) \geq b(M/g \setminus e)$ . Thus  $b(M/f/e) \leq b(M/g/e)$ . Since  $f$  is freer than  $g$  in  $M/e$ , it follows, by Corollary 8(i), that



$b(M/f/e) = b(M/g/e)$ . Consequently,  $b(M/f \setminus e) = b(M/g \setminus e)$ . Therefore, by the induction assumption,  $f$  and  $g$  are clones in  $M \setminus e$ .

Since  $f$  and  $g$  are not clones in  $M$ , there is a circuit  $C$  of  $M$  containing  $g$  such that  $f \notin \text{cl}_M(C)$ . Assume there is an element  $e \in E(M) - (C \cup f)$ . Then  $C$  is a circuit of  $M \setminus e$  containing  $g$  such that  $f \notin \text{cl}_{M \setminus e}(C)$ . Hence  $f$  and  $g$  are not clones in  $M \setminus e$ , a contradiction. It follows that  $C = E(M) - \{f\}$ . Thus  $r_M(C) = r(M \setminus f) = r(M) - 1$ . Therefore  $f$  is a coloop of  $M$ , a contradiction.  $\square$

**Proposition 12.** *Let  $f$  be freer than  $g$  in  $M$ . Let  $L$  be obtained from  $M$  by deleting every element of  $E(M) - \{f, g\}$  that is parallel to  $g$ . Then  $h(f; M) = h(g; M)$  if and only if  $f$  and  $g$  are clones in  $L$ .*

*Proof.* Suppose  $f$  and  $g$  are clones in  $L$ . As  $h(f; M) = h(f; L)$  and  $h(g; M) = h(g; L)$ , we have  $h(f; M) = h(g; M)$ . To prove the converse, suppose that  $h(f; M) = h(g; M)$ . Let  $\mathcal{H}(M; f; \bar{g})$  be the set of hyperplanes of  $M$  containing  $f$  but not  $g$ . Then  $|\mathcal{H}(M; f; \bar{g})| = |\mathcal{H}(M; g; \bar{f})|$ . Hence  $|\mathcal{H}(L; f; \bar{g})| = |\mathcal{H}(L; g; \bar{f})|$ . Clearly, if  $\{f, g\}$  is a 2-circuit of  $L$ , then  $f$  and  $g$  are clones in  $L$ . Thus, we may assume that  $g$  is not any 2-circuit of  $L$ .

For  $J \in \mathcal{H}(L; g; \bar{f})$ , let  $B_J$  be an arbitrarily chosen basis of  $J$  containing  $g$ . Then  $\text{cl}_L(B_J - g)$  is a rank- $(r-2)$  flat of  $L$ . Let  $\text{cl}_L(B_J - g) \cup f = F$ . Then  $r(F) = r - 1$  otherwise  $f \in \text{cl}_L(B_J - g)$ , so  $g \in \text{cl}_L(B_J - g)$ , a contradiction. Consider  $\text{cl}_L(F)$ . Assume  $f$  is not a coloop of  $\text{cl}_L(F)$ . Then  $\text{cl}_L(F)$  contains a circuit  $C$  containing  $f$ . Since  $f$  is freer than  $g$ , we see that  $g \in \text{cl}_L(F)$ . Then  $\text{cl}_L(F) \supseteq B_J$ . Thus  $\text{cl}_L(F) \supseteq J$ . As  $r(\text{cl}_L(F)) = r(J)$ , we deduce that  $\text{cl}_L(F) = J$ . But  $f \notin J$ , a contradiction. Thus  $f$  is a coloop of  $\text{cl}_L(F)$ , so  $\text{cl}_L(B_J - g) \cup f \in \mathcal{H}(L; f; \bar{g})$ . Let  $\psi(J) = \text{cl}_L(B_J - g) \cup f$ . Note that  $\psi$  depends upon the choices made for the bases  $B_J$ . Moreover,  $\psi$  maps  $\mathcal{H}(L; g; \bar{f})$  to  $\mathcal{H}(L; f; \bar{g})$ .

**12.1.**  $\psi$  is bijective.

To see that  $\psi$  is injective, suppose that, for distinct members  $J_1$  and  $J_2$  of  $\mathcal{H}(L; g; \bar{f})$ , the hyperplanes  $\psi(J_1)$  and  $\psi(J_2)$  are equal. Then  $\text{cl}_L(B_{J_1} - g) = \text{cl}_L(B_{J_2} - g)$ . Now the rank- $(r-2)$  flat  $J_1 \cap J_2$  contains  $g$  and so contains the rank- $(r-1)$  set  $B_{J_1}$ , a contradiction. Since  $|\mathcal{H}(L; g; \bar{f})| = |\mathcal{H}(L; f; \bar{g})|$  and  $\psi$  is injective, we conclude that  $\psi$  is bijective.

**12.2.**  $g$  is a coloop of  $L|J$  for every  $J \in \mathcal{H}(L; g; \bar{f})$ .

Suppose  $g$  is not a coloop of  $L|J$ . Then  $\text{cl}_L(B_J - g)$  is a subset of  $J$  avoiding  $g$ . As  $g$  is not a coloop, there is an element  $h$  of  $J - \text{cl}_L(B_J - g)$ . Since  $g$  is not in any 2-circuits of  $L$ , the elements

$g$  and  $h$  are not parallel. Thus  $\{g, h\}$  is independent. Extend  $\{g, h\}$  to a basis  $B'_J$  of  $L|J$ . Then  $\text{cl}_L(B'_J - g) \neq \text{cl}_L(B_J - g)$  because  $h \in \text{cl}_L(B'_J - g) - \text{cl}_L(B_J - g)$ . Thus  $\text{cl}_L(B'_J - g) \cup f$  is a member of  $\mathcal{H}(L; f; \bar{g})$  that is not in the set  $\psi(\mathcal{H}(L; g; \bar{f}))$ . As  $|\mathcal{H}(L; g; \bar{f})| = |\mathcal{H}(L; f; \bar{g})|$  and  $\psi$  is a bijection, this is a contradiction.

Suppose  $g$  is not freer than  $f$  in  $L$ . Then  $L$  has a cyclic flat  $K$  containing  $g$  and avoiding  $f$ . Take a basis  $B$  for  $K$  and consider  $B \cup f$ . Extend  $B \cup f$  to get a basis  $B_L$  for  $L$ . Then  $\text{cl}_L(B_L - f)$  is a hyperplane of  $L$  containing  $g$  and avoiding  $f$ . Moreover, since  $K \subseteq \text{cl}_L(B_L - f)$ , the hyperplane  $\text{cl}_L(B_L - f)$  has a circuit containing  $g$ ; that is,  $g$  is not a coloop of  $\text{cl}_L(B_L - f)$ , a contradiction to 12.2.  $\square$

The next corollary is obtained by applying Proposition 12 to  $M^*$ .

**Corollary 13.** *Let  $f$  be freer than  $g$  in  $M$ . Let  $N$  be obtained from  $M$  by contracting every element of  $E(M) - \{f, g\}$  that is in series with  $f$ . Then  $\gamma(f; M) = \gamma(g; M)$  if and only if  $f$  and  $g$  are clones in  $N$ .*

*Proof.* Clearly  $\gamma(f; N) = \gamma(g; N)$  if  $f$  and  $g$  are clones in  $N$ . Hence  $\gamma(f; M) = \gamma(g; M)$ . To prove the converse, suppose that  $\gamma(f; M) = \gamma(g; M)$ . Then  $\gamma(f; N) = \gamma(g; N)$ . If  $f$  and  $g$  are in series in  $N$ , then  $f$  and  $g$  are clones in  $N$ . Thus, we will assume that  $f$  and  $g$  are not a series pair in  $N$ .

Let  $\mathcal{C}(M; f, \bar{g})$  be the set of circuits of  $M$  containing  $f$  and avoiding  $g$ . Then  $|\mathcal{C}(N; f; \bar{g})| = |\mathcal{C}(N; g; \bar{f})|$ . By duality,  $|\mathcal{C}(N; f, \bar{g})| = |\mathcal{H}(N^*; g; \bar{f})|$ . Therefore  $|\mathcal{H}(N^*; g; \bar{f})| = |\mathcal{H}(N^*; f; \bar{g})|$ , and, consequently,  $h(g; M^*) = h(f; M^*)$ . Since  $g$  is freer than  $f$  in  $M^*$ , by Proposition 12, we see that  $f$  and  $g$  are clones in  $M^* \setminus X$  where  $X$  is the set of elements of  $E(M) - \{f, g\}$  that are parallel to  $f$  in  $M^*$ . It follows that  $f$  and  $g$  are clones in  $N$ .  $\square$

Based on Theorem 11, Proposition 12, and Corollary 13, one may guess that, when  $f$  is freer than  $g$  in  $M$  and  $\gamma'(f; M) = \gamma'(g; M)$ , the elements  $f$  and  $g$  must be clones in  $M$  when  $f$  is not in any 2-cocircuit of  $M$ . To see that this is not so, let  $M$  be the rank-5 matroid that is obtained by taking the 2-sum across a common basepoint  $p$  of two 4-point lines  $M_2$  and  $M_3$  and of a 6-element rank-3 matroid  $M_1$  that has  $\{g, a, b\}$  as its only non-spanning circuit and has  $f, p$ , and  $c$  as free elements. Then  $f$  is freer than  $g$  in  $M$  and  $\gamma'(f; M) = 0 = \gamma'(g; M)$ . But  $f$  and  $g$  are not clones in the cosimple matroid  $M$ .

The *truncation* of  $M$ , which we will denote  $\tau(M)$ , is the matroid obtained from  $M$  by taking the free extension  $M +_{E(M)} e$  of  $M$  by  $e$  and then contracting the free element  $e$ . Note that we use  $\tau(M)$  rather

than the more standard  $T(M)$  to denote truncation in order to avoid confusion with the Tutte polynomial of  $M$ .

**Corollary 14.** *Let  $r(M) = r \geq 2$ . If  $f$  is freer than  $g$  in  $M$ , then  $W_{r-2}(f; M) = W_{r-2}(g; M)$  if and only if  $f$  and  $g$  are clones in the matroid obtained from  $\tau(M)$  by deleting every element of  $E(M) - \{f, g\}$  that is parallel to  $g$ .*

*Proof.* Let  $F$  be a rank- $(r-2)$  flat of  $M$ . Then  $F$  is a rank- $(r-2)$  flat of  $M +_{E(M)} e$  avoiding  $e$ . Hence  $F$  is a hyperplane of  $\tau(M)$ . Therefore, the rank- $(r-2)$  flats of  $M$  containing an element  $x$  are precisely the hyperplanes of  $\tau(M)$  containing  $x$ , that is,  $W_{r-2}(f; M) = W_{r-2}(g; M)$  if and only if  $h(f; \tau(M)) = h(g; \tau(M))$ . As  $f$  is freer than  $g$  in  $\tau(M)$ , the result follows immediately from Proposition 12.  $\square$

The following result generalizes Corollary 14 to the  $i$ -th truncation  $\tau^i(M)$  of  $M$ , defined recursively by  $\tau^i(M) = \tau(\tau^{i-1}(M))$  where  $\tau^0(M) = M$ .

**Corollary 15.** *Suppose  $f$  is freer than  $g$  in  $M$  and  $r(M) = r \geq 2$ . Then  $W_k(f; M) = W_k(g; M)$  for some  $k$  with  $1 \leq k \leq r-1$  if and only if  $f$  and  $g$  are clones in the matroid obtained from  $\tau^{r-k-1}(M)$  by deleting every element of  $E(M) - \{f, g\}$  that is parallel to  $g$ .*

*Proof.* This follows immediately from Corollary 14.  $\square$

## REFERENCES

- [1] T. Brylawski and J. Oxley, The Tutte polynomial and its applications, *Matroid Applications*, N. White, Ed., Cambridge University Press, Cambridge, pp. 123–225, 1992.
- [2] R. Duke, Freedom in matroids, *Ars Combin.* **26B** (1988), 191–216.
- [3] J. Geelen, J. Oxley, D. Vertigan, and G. Whittle, Totally free expansions of matroids, *J. Combin. Theory Ser. B* **84** (2002), 130–179.
- [4] F. Jaeger, D. L. Vertigan, and D. J. A. Welsh, On the computational complexity of the Jones and Tutte polynomials, *Math. Proc. Cambridge Phil. Soc.* **108** (1990), 35–53.
- [5] D. Lucas, Weak maps of combinatorial geometries, *Trans. Amer. Math. Soc.* **206** (1975), 247–280.
- [6] J. Oxley, *Matroid Theory*, Second Edition, Oxford University Press, New York, 2011.
- [7] J. P. S. Kung and H. Q. Nguyen, Weak maps, *Theory of Matroids*, N. White, Ed., Cambridge University Press, Cambridge, pp. 254–271, 1986.

MATHEMATICS DEPARTMENT, LOUISIANA STATE UNIVERSITY, BATON ROUGE,  
LOUISIANA

*Email address:* ccho3@lsu.edu